

Inhibition of quorum sensing virulente factors of *Pseudomonas aeruginosa* PAO1 by *Ferulago macedonica* and *Echinophora sibthorpiana* extracts and essential oils

KSENIJA MILESKI^{1,*}, ANA ĆIRIĆ², VLADO MATEVSKI^{3,4}, PETAR MARIN¹, MARINA SOKOVIĆ², AND ANA DŽAMIĆ¹

¹Institute of Botany and Botanical Garden "Jevremovac", Faculty of Biology, University of Belgrade, Studentski trg 16, 11000 Belgrade, Serbia

²Dep. of Plant Physiology, Institute for Biological Research "Siniša Stanković", University of Belgrade, Bul. Despota Stefana, 142, 11000 Belgrade, Serbia

³Institute of Biology, Faculty of Natural Sciences and Mathematics, University "S. Kiril and Metodij", Gazi Baba, 1000 Skopje, Macedonia

⁴Macedonian Academy of Sciences and Arts, Bul. Krste Misirkov, 2, P.O. Box 428, 1000 Skopje, Macedonia

*Corresponding author: ksenija.mileski@bio.bg.ac.rs

Received: December 8, 2017

Accepted: December 13, 2017

Published on-line: December 14, 2017

Published: December 25, 2017

This study was conducted to analyse extracts and essential oils of *Ferulago macedonica* and *Echinophora sibthorpiana* (Apiaceae) for anti-quorum sensing potential using *Pseudomonas aeruginosa* PAO1 as biomonitoring system. Anti-quorum sensing screening was carried out by three different tests which showed that examined samples possessed significant effects in suppressing of some virulent determinants of PAO1. The essential oils of *F. macedonica* and *E. sibthorpiana* aerial parts reduced the growth and modified the appearance of *P. aeruginosa* colonies and flagella to a large extent in twitching and motility assay. In the presence of essential oil of *E. sibthorpiana*, diameter of the colony was 6.67 mm, the flagella were absent and the color of the colony was changed from green to white. In anti-biofilm assay, the inhibitory activity of *E. sibthorpiana* was more notable than of *F. macedonica*. Applied colorimetric test revealed that examined samples strongly reduced the production of pigment pyocyanin where the highest inhibition of its synthesis showed *F. macedonica* oil (5.22%), followed by *E. sibthorpiana* oil (10.69%), which was significantly higher potential in comparison to used antibiotics (84.27% and 97.59% for streptomycin and ampicillin, respectively).

Key words: Apiaceae, *Ferulago macedonica*, *Echinophora sibthorpiana*, anti-quorum sensing, *Pseudomonas aeruginosa* PAO1

<http://dx.doi.org/10.5937/leksi1737033M>

1. INTRODUCTION

Studied species *Ferulago macedonica* Micevski et Mayer and *Echinophora sibthorpiana* (Guss.) Tutin are herbaceous perennial plants which belong to the carrot family (Apiaceae). Endemic *F. macedonica*, with upright stem and yellow inflorescence, can be found in a limestone area of Republic of Macedonia. It belongs to the genus *Ferulago* W. Koch which includes around 50 species spread in Iran, Turkestan and in surrounding of Adriatic Sea (Canonn, 1968). Genus *Echinophora* L. consists of species *E. tenuifolia* L. and *E. spinosa* L. which are distributed from Mediterranean region to Crete and Crimea. In flora Europaea it is recorded that species *E. tenuifolia* consists of subsp. *tenuifolia* and subsp. *sibthorpiana* (Tutin, 1968). According to recent data, *E. tenuifolia* subsp. *sibthorpiana* is treated as separate species *E. sibthorpiana* (Micevski, 2005). This species has strong roots, hairy leaves, branched stem and yellow inflorescence. It

inhabits the mountain meadows and saline areas of Republic of Macedonia (Micevski, 2005).

Many Apiaceae species are known as aromatic and medicinal plants with wide usage in human diet, as well as in traditional and official medicine. Some species of *Ferulago* and *Echinophora* genera are utilized since ancient times as a flavoring and digestive agents. Thus, certain *Ferulago* species are added as spices to dishes or to melted fats for aromatization and as food preservatives. A few representatives of this taxa are used in traditional medicine as sedatives, tonics and in the treatment of intestinal wounds due to their antiparasitic effect. Moreover, they are used against ulcers, snake bites, headache and diseases of the spleen (Khalighi-Sigaroodi et al., 2005). In Turkey and Iran, extracts of the roots of *Ferulago* species are applied as aphrodisiacs. Also, their fragrance is used in the perfumery and cosmetic industry (Taran et al., 2010). Tarana

herb which is local name for *E. sibthorpiana* in Turkey gives pleasant aroma and flavor in preparation of tarana—one of the oldest traditional Turkish dishes made of cereals. It was proven that this species stimulates the fermentation of tarana by affecting lactic acid bacteria and yeast *Saccharomyces cerevisiae*. In traditional medicine, it is used as a fungicide for the treatment of wounds and stomach ulcers, also, as digestive and anti-spasm agent (Baser et al., 1998; Cakilcioglu and Turkoglu, 2010; Değirmencioglu et al., 2005; Gurbuz et al., 2010; Ozdemir et al., 2007). In addition, the seeds and the roots of this taxon have been effectively applied in the treatment of epilepsy (Eadie, 2004).

Previous studies on chemical analyses of *F. macedonica* and *E. sibthorpiana* showed that they possess various secondary metabolites. Thus, the chemical composition of *Ferulago* species revealed that they contain coumarins, polyacetylenes, flavonoids, monoterpenes, sesquiterpenes and aromatic compounds such as phenylpropanoids (Demirci and Özkan, 2014; Razavi et al., 2015; Ruberto et al., 1994). Regarding chemical analyses of *Echinophora* species, essential oils were studied most frequently. In essential oil of aerial parts of *E. sibthorpiana*, α -phellandrene and methyl-eugenole were the most abundant components (Ahmad et al., 1999; Baser et al., 1994; 1998; Gokbulut et al., 2013). In extracts obtained from aerial parts of *E. platyloba*, stigmasterol, sitosterol, and stigmasterol- β -D-glycoside were identified as dominant constituents (Valizadeh et al., 2014). In regard to the complex chemical composition, *Ferulago* species exhibit numerous biological activities. The antimicrobial aspect of essential oils of *Ferulago* species was examined in few previous works (Cecchini et al., 2010; Ozkan et al., 2008; Taran et al., 2010). Essential oil of *F. campestris* showed to be a promising anti-*Candida* agent with extremely strong activity on clinically isolated strains of *C. albicans*, *C. tropicalis*, and *C. glabrata*, thus corroborating the historical usage of this plant in the treatment of candidiasis (Cecchini et al., 2010). The various extracts and essential oils of *Echinophora* species *in vitro* also showed diverse biological activities and the most prominent were their antioxidant (Gholivand et al., 2011; Gokbulut et al., 2013; Saei-Dehkordi et al., 2012) and antimicrobial potentials (Avijgan et al., 2010; 2012; Delaram et al., 2011; Entezari et al., 2009; Sharafati-chaeshtori et al., 2012; Yousef et al., 2012). Moreover, it was found that flower, leaf, and stem essential oils from *Echinophora lamondiana* and from aerial parts of *Echinophora tenuifolia* ssp. *sibthorpiana* act as biopesticides and repellents (Ali et al., 2015; Evergetis et al., 2013).

This work represents the extension of our previous research, where *F. macedonica* and *E. sibthorpiana* were characterized for chemical profiles of essential oils and extracts. The most abundant compounds in essential oils of *F. macedonica* inflorescence and aerial parts were α -pinene (43.1% and 22.8%, respectively) and sabinene (26.7% and 15.5%, respectively), while total phenolic and flavonoid contents in all extracts ranged between 34.12-59.68 mg GAE/g dw and 9.1-23.7 mg QE/g dw, respectively (Mileski et al., 2015). In the essential oil of *E. sibthorpiana* methyl eugenol (60.4%), p-cymene (11.2%) and α -phellandrene (10.2%) were main compounds, while total phenolic and flavonoid concentrations in the extracts ranged from 38.6-60.7 mg GAE/g dw, and 3.1-19.0 mg QE/g dw, respectively (Mileski et al., 2014). The aim of this study was to determine the anti-quorum sensing (anti-QS) activity of *F. macedonica* and *E. sibthorpiana* on *P. aeruginosa* PAO1 taking into consideration the widespread use of related representatives as antimicrobial agents in traditional medicine. This analysis was conducted by determining the ability of extracts and essential oils to reduce twitching and flagella motility, biofilm formation and pyocyanin production of this bacterium and the comparison of their activity with the effects of commercial

antibiotics was done.

2. MATERIALS AND METHODS

1. Chemicals

For antimicrobial assays, several solvents, culture broth, and standards were used. Organic solvents were purchased from „Zorka pharma“, Šabac, Serbia. Tryptic Soy Broth (TSB) was obtained from the Institute of Immunology and Virology „Torlak“, Belgrade, Serbia, while streptomycin and ampicillin solutions (1 mg/mL) from Hyclone, Logan, Ut, USA. Dimethylsulfoxide (DMSO) was purchased from Merck KGaA, Germany and phosphate-buffered saline (PBS) were from Sigma Chemical Co., St. Louis, USA.

2. Plant Material

Both species were collected in flowering stage at their natural sites of the Republic of Macedonia and determined by one of the authors, Prof. V. S. Matevski. Plant material of *F. macedonica* was collected around the Negotino city (GPS: N 41°30,362', E 22°06,091'), while *E. sibthorpiana* was collected around the Štip city (GPS: N 41°50,324'; E 22°08,289') in July 2012. Voucher specimens of *F. macedonica* (BEOU 16659) and *E. sibthorpiana* (BEOU 16658) are deposited at the Herbarium of the Institute of Botany and Botanical Garden „Jevremovac“, (BEOU), Faculty of Biology, University of Belgrade, Serbia.

3. Plant extraction

Methanol, ethanol and aqueous extracts of *F. macedonica* (inflorescence and aerial parts) and *E. sibthorpiana* (aerial parts and roots) were obtained by ultrasonic extraction procedure. Powdered plant materials (10 g) were extracted for 24 h in the dark with 200 mL of listed solvents using an ultrasonic apparatus for the first and the last hour of extraction. After filtration, methanol and ethanol were evaporated under reduced pressure at 40°C, while frozen aqueous were lyophilized. Measured extracts (Table 1) were kept in the glass bottles at 4°C until further application.

Table 1. Obtained yields of *Ferulago macedonica* and *Echinophora sibthorpiana* crude extracts.

Species	Extract	Plant part	Yield [g]
<i>F. macedonica</i>	Methanol	Inflorescence	3.751
		Aerial parts	1.332
	Ethanol	Inflorescence	2.157
		Aerial parts	1.179
Aqueous	Inflorescence	2.020	
	Aerial parts	1.351	
<i>E. sibthorpiana</i>	Methanol	Aerial parts	0.841
		Roots	1.049
	Ethanol	Aerial parts	0.794
		Roots	0.482
	Aqueous	Aerial parts	0.974
		Roots	1.284

4. Essential oil isolation

Yellowish essential oils of *F. macedonica* and *E. sibthorpiana* were isolated by hydrodistillation (3h) from 200 g of dry plant material using Clevenger apparatus (Ph.Eur.8.0., 2013). Obtained essential oils from aerial parts of *F. macedonica* (without inflorescence) and herbal parts of *E. sibthorpiana* (the whole plant) were refrigerated in a sealed vial prior to anti-QS analyses (0.61% and 0.43%, w/w-dry bases, respectively).

5. Determination of anti-QS activity

5.1. Preparation of extracts

Crude extracts of studied species were dissolved in 5% DMSO to obtain the stock concentrations of 20 mg/mL for *F. macedonica* and 30 mg/mL for *E. sibthorpiana* samples. Subsequently, obtained solutions were diluted in appropriate ratio for further analyses.

5.2. Bacterial strain, growth media and culture conditions

The bacterium *Pseudomonas aeruginosa* PAO1 from Institute for Biological Research "Siniša Stanković" in Belgrade, Serbia was used to perform the experiments. *P. aeruginosa* was routinely grown in Luria-Bertani medium (1% w/v NaCl, 1% w/v Tryptone, 0.5% w/v yeast extract) with shaking (220 rpm) and cultured at 37°C.

5.3. Microwell dilution assay

MIC values for *P. aeruginosa* were determined by microwell dilution method following the protocol described by CLSI (2009) and Hanel and Raether (1998). The technique was carried out using pure, undiluted essential oils and different dilutions of extracts in 5% DMSO (stock concentrations were 20 mg/mL and 30 mg/mL for *F. macedonica* and *E. sibthorpiana*, respectively). For detection of bacterial growth, a colorimetric viability assay was used (CLSI, 2009; Tsukatani et al. 2012). Obtained results were compared with positive control (solution of streptomycin in the concentration of 1 mg/mL).

5.4. Biofilm formation

To determinate the influence of tested samples on biofilm formation of *P. aeruginosa*, the procedure of Drenkard and Ausubel (2002) and Spoering and Lewis (2001) with slight modifications was followed. This test was applied to investigate the ability of used extracts and essential oils to reduce/stop the biofilm formation in different sub-inhibitory concentrations (subMICs) (0.125, 0.25, 0.5 of MICs) which were determined in our previous studies (Mileski et al., 2014; 2015). The experiment was done in polystyrene flat-bottomed microtitre 96-well plates. In brief, 100 µL of the medium (control) was pipetted into each well of the plates. Subsequently, 0.1 mL of subMICs of tested samples and 100 µL of overnight culture inoculum was added. After 24 h of incubation at 37°C, each well was washed twice with sterile PBS (pH 7.4) and dried. Afterward, 0.1% of crystal violet was poured and left for 10 min with the aim to determine formed biofilm mass. After drying, 0.2 mL of 95% ethanol (v/v) was added to solubilize the dye that had stained the biofilm cells. The excess stain was washed off with distilled water and left for 10 min. After homogenization of the contents of the wells, the absorbance was read at $\lambda=625$ nm on a Sunrise™-Tecan ELISA reader.

Biofilm formation values were calculated using formula:

$$\text{Biofilm formation [\%]} = \frac{\text{mean } A_{620} \text{ control well}}{\text{mean } A_{620} \text{ treated well}} \times 100\%$$

The experiment was done in triplicate. The values are presented as mean values \pm standard deviations.

5.5. Twitching and flagella motility

The cells of *P. aeruginosa* grown in the presence or the absence of 0.5 MICs of the extracts and essential oils were washed twice and re-suspended in sterile PBS to gain 1×10^8 CFU/mL (optical density (OD) of 0.1 at 660 nm). The cells were stabbed into a nutrient agar plate with a sterile toothpick and incubated overnight at 37°C and later incubated at room temperature for 48 h. The zone of the motility and the colony edges were observed and measured with a light microscope (Leika, 020–518.500 DM LS, Germany) (O'Toole and Kolter, 1998b;a). For testing flagella motility, all samples were mixed into 10

mL of molten MH and poured immediately over the surface of a solidified LB agar plate as an overlay. The plate was point inoculated with an overnight culture of PAO1 once the overlaid agar had solidified and incubated at 37°C for 3 days. The extent of swimming was determined by measuring the diameters of the colony in three different directions with square millimeter using graph paper (Yeo and Tham, 2012). The experiment was done in triplicate and the values are presented as a means \pm standard deviations.

5.6. Pyocyanin production

The flask assay was used for the quantification of the ability of tested samples (0.5 MICs) to inhibit pyocyanin production in *P. aeruginosa*. An overnight culture of PAO1 was diluted to OD₆₀₀ 0.2. SubMICs of the samples were added to 5 mL of bacterial inocula and the incubation lasted for 24 h at 37°C. Thereafter, the treated cultures were extracted with 3 mL of chloroform, followed by mixing the chloroform layer with 1 mL of 0.2 M HCl. The absorbance readings of the extracted organic layer were conducted at $\lambda=520$ nm using a Shimadzu UV1601 spectrophotometer (Kyoto, Japan) (Yeo and Tham, 2012). The experiment was done in triplicate. The values are expressed as a ratio (OD₅₂₀/OD₆₀₀) \times 100.

5.7. Statistical analysis

The results of this study were analyzed using one-way analysis of variance (ANOVA) followed by Tukey's HSD test with $P < 0.05$ assumed as significant difference among mean values. This analysis was carried out using SPSS v. 18.0 program.

3. RESULTS AND DISCUSSION

1. Anti-QS activity of extracts and essential oils

The main goal of this study was to determinate the anti-QS activity of tested samples on *P. aeruginosa* PAO1. The results obtained in the analyses are given in Tables 2 and 3 and Figures 1-3. There is an increasing number of evidence that plants produce anti-QS compounds that interfere with intercellular signaling communication (Bauer and Mathesius, 2004; Teplitzki et al., 2000). So far, several potential mechanisms of actions for anti-QS agents have been proposed, which are mainly based on interfering the QS system. They include inhibition of genetic regulatory system or biosynthesis of autoinductors, their inactivation or degradation and the reaction with signal receptors (Glamočlija et al., 2015). In recent years, the representatives of the Apiaceae family are also studied for QS inhibition in the bacteria. So far, only a few species were explored for QS potential (Khan et al., 2009; Sepahi et al., 2015) and this is the first data on *F. macedonica* and *E. sibthorpiana* activity against QS communication in *P. aeruginosa*. In our earlier investigations, the antibacterial potential of tested species and MIC values for *P. aeruginosa* were defined and the essential oils of *E. sibthorpiana* and *F. macedonica* showed to be the strongest anti-*Pseudomonas* agents among tested samples, in the range of streptomycin (MICs=8.00–10.00 mg/mL and 1.50–12.00 mg/mL for extracts; MICs=0.60 mg/mL and 0.34 mg/mL for essential oils of *F. macedonica* and *E. sibthorpiana*, respectively) (Mileski et al., 2014; 2015). According to presented results, it was observed that QS profiles of used samples differed significantly and used subMICs exhibited inhibitory activity on PAO1 determinants in varying quantity. Thus, comparative analysis of the results obtained in twitching and flagella test showed that in the presence of some samples, *P. aeruginosa* colonies were altered and the flagella appearance and diameters were affected. Control colony of this bacterium grew rapidly due to the ability of the cells to move using flagella with regular size, creating a wide zone of 12.00 mm in diameter, with rough, uneven and irregular edges. Bacterial

Table 2. Effect of *Ferulago macedonica* and *Echinophora sibthorpiana* extracts and essential oils on PAO1 twitching and flagella motility.

Extract/Standard	Plant part	Colony diameter ^a [mm]	Flagella diameter [μm]	Colony color	Colony edge
<i>F. macedonica</i>					
Methanolic	Inflorescence	14.33 ± 1.15 c	–	Light green	–
	Aerial parts	10.33 ± 2.08 ab	64 – 160	Transparent white	Sporadical flagella
Ethanollic	Inflorescence	13.66 ± 1.53 b	40 – 240	Pale brown	Regular flagella
	Aerial parts	15.33 ± 1.53 c	80 – 240	Transparent white	Regular flagella
Aqueous	Inflorescence	14.66 ± 0.58 c	40 – 160	Whitish	Reduced flagella
	Aerial parts	16.33 ± 1.00 c	40 – 160	Pale brown	Regular flagella
Ess. oil	Aerial parts	7.33 ± 2.08 a	16 – 40	Whitish	Rare flagella
<i>E. sibthorpiana</i>					
Methanolic	Aerial parts	14.00 ± 4.58 c	32 – 120	White	Rare flagella
	Roots	15.00 ± 2.00 c	24 – 120	Transparent white	Very rare flagella
Ethanollic	Aerial parts	15.33 ± 1.53 c	40 – 120	Whitish	Rare flagella
	Roots	14.00 ± 0.00 c	80 – 160	Pale brown	Regular flagella
Aqueous	Aerial parts	15.66 ± 0.58 c	56 – 136	Whitish	Reduced flagella
	Roots	14.00 ± 1.00 c	40 – 120	Pale brown	Reduced flagella
Ess. oil	Aerial parts	6.67 ± 0.51 a	–	Whitish	–
Streptomycin		11.00 ± 1.00 b	24 – 56	Green	Tiny flagella
Ampicillin		13.33 ± 5.03 bc	16 – 56	Green	Regular flagella
Control (10 ⁹ CFU/mL)		12.00 ± 1.00 b	56 – 80	Light green	Regular flagella

^aValues with different indicated letters in the same column mean significant difference (P<0.05).

cells formed thin layers with regular surface and blurred, unclear zones around green colony (Figures 1 and 2). According to the results, the samples reduced the movement of the cells and the growth of the colonies. As it can be seen in Table 2, the colors of *P. aeruginosa* treated with *F. macedonica* extracts and essential oil differed from transparent white to pale brown. Regarding *F. macedonica*, the best reduction of all parameters in this assay showed the essential oil of aerial parts in which presence the colony diameter was 7.33 mm (Table 2 and Figure 1). Moreover, it has been shown that both, essential oil and methanol extract of inflorescence had stronger activity in comparison to ampicillin (small diameters, absence of flagella), while methanol extract of aerial parts showed better activity than streptomycin. In this examination, the extracts and essential oil of *E. sibthorpiana* showed the following results: the color of the colonies ranged from transparent white, through whitish to pale brown (Table 2). The diameters of the colonies that grew in the presence of *E. sibthorpiana* extracts were greater in comparison to the control (14.00–15.66 mm and 12.00 mm, respectively) and colonies with antibiotics (11.00–13.00 mm). The essential oil of *E. sibthorpiana* aerial parts showed the best activity on all used parameters (Table 2, Figure 2). Thus, in the presence of the oil, the diameter of the whitish colony was 6.67 mm and flagella were not detected. It can be concluded that only essential oil of this species showed stronger activity than synthetic antibiotics. In this assay, few extracts did not show inhibitory effect on *P. aeruginosa* motility since the color and appearance of the colonies and flagella remained unchanged (Table 2; Figures 1 and 2).

The effects of used extracts and essential oils on PAO1 biofilm formation are presented in Table 3. Samples were tested at concentrations of 0.5, 0.25 and 0.125 of MICs. In general, anti-biofilm activity of *E. sibthorpiana* was more notable than of *F. macedonica*. The results given in Table 3 indicate low activity or complete absence of the effect of some *F. macedonica* samples in the reduction of biofilm formation. The extent of reducing ability ranged from 2.93% for 0.125 MIC of aerial part aqueous extract to 28.22% for 0.5 MIC of *F. macedonica* essential oil. The best activity among extracts showed 0.25 MIC

of ethanol extract of the inflorescence. On the other hand, both methanol extracts did not exhibit effectiveness in suppressing of biofilm formation. It can be concluded that streptomycin and ampicillin had stronger activity than extracts and essential oil of this species (Table 3). According to the results obtained for *E. sibthorpiana*, only methanol and aqueous extracts of roots showed activity at all tested concentrations (Table 3). The largest reduction of biofilm production was measured in the presence of 0.5 MIC of root methanol extract (59.65%), while 0.125 MIC of this sample reduced only 0.29% of its synthesis. In comparison with streptomycin and ampicillin, only methanol extract of the roots had stronger effect on biofilm (0.5 MIC), while ethanol extracts did not inhibit the biofilm synthesis (Table 3).

The reducing effects of tested agents on pyocyanin production were monitored by colorimetric test and it was conducted through comparison of the intensity of green coloration of the samples and control. The applied amounts of extracts and essential oils were sufficient to reduce the concentration of synthesized pigment and recorded inhibitory capacities of analyzed species in this assay were the most prominent (Figure 3). According to the results, essential oil of *F. macedonica* showed the strongest inhibition of pyocyanin synthesis (5.22%), followed by *E. sibthorpiana* oil (10.69%). Among extracts, the most effective were ethanol extract of *E. sibthorpiana* roots (14.11%) and of *F. macedonica* aerial parts (19.05%), which were significantly lower concentrations of synthesized pigment in comparison to the control (141.55%). SubMICs of *F. macedonica* showed much higher activity in the reduction of pigment's production than in biofilm synthesis. The best *F. macedonica* extract in inhibition of pyocyanin production was ethanol extract of aerial parts (19.05%). The highest measured concentration of the pigment was with methanol extract of aerial parts (68.6%) which was still lower amount than in the presence of streptomycin and ampicillin (84.27% and 97.59%, respectively). In addition, all tested *E. sibthorpiana* samples significantly reduced the synthesis of pyocyanin and based on the reduction level it can be concluded that *E. sibthorpiana* showed strong anti-pyocyanin activity. The essential oil of

Table 3. Effect of *Ferulago macedonica* and *Echinophora sibthorpiana* extracts and essential oils on PAO1 biofilm formation

Extract/Standard	Plant part	0.5 MIC ^{a,b} [%]	0.25 MIC [%]	0.125 MIC [%]
<i>F. macedonica</i>				
Methanolic	Inflorescence	n.i.	n.i.	n.i.
	Aerial parts	n.i.	n.i.	n.i.
Ethanollic	Inflorescence	n.i.	n.i.	86.70 ± 0.00 b
	Aerial parts	n.i.	96.18 ± 0.78 b	81.60 ± 1.18 a
Aqueous	Inflorescence	n.i.	76.17 ± 1.21 ab	80.41 ± 0.95 a
	Aerial parts	n.i.	n.i.	97.07 ± 0.53 b
Ess. Oil	Aerial parts	71.78 ± 2.04 b	87.72 ± 1.08 b	n.i.
<i>E. sibthorpiana</i>				
Methanolic	Aerial parts	75.15 ± 1.35 b	n.i.	n.i.
	Roots	40.35 ± 2.51 a	66.52 ± 2.03 a	99.71 ± 0.82 b
Ethanollic	Aerial parts	n.i.	n.i.	n.i.
	Roots	n.i.	n.i.	n.i.
Aqueous	Aerial parts	92.11 ± 0.88 b	n.i.	n.i.
	Roots	86.84 ± 1.65 b	71.05 ± 1.72 a	77.37 ± 1.45 a
Ess. Oil	Aerial parts	n.i.	n.i.	68.42 ± 2.20 a
Streptomycin		69.16 ± 0.65 b	56.46 ± 0.46 a	92.16 ± 0.37 b
Ampicillin		49.40 ± 0.46 a	70.97 ± 0.36 a	88.36 ± 0.42 b

^aThe values are presented as mean values ± standard deviations; n.i. - not identified.

^bValues with different indicated letters in the same column mean significant difference (P<0.05).




Fig. 1. Light microscopy of colony edges of PAO1 in twitching motility, grown in the presence or absence of *Ferulago macedonica* extracts, essential oil and antibiotics. The bacterial colonies grown with the presence of 0.5 MIC of extracts (A-F) and essential oil (G); *Pseudomonas aeruginosa* colony in the presence of streptomycin had reduced protrusion (H); colony in the presence of ampicillin with regularly formed protrusions (I); control *Pseudomonas aeruginosa* produced a flat, widely spread, irregularly shaped colony (J); Magnification: (A-J) × 100.

this species had the highest potency reducing 130.86% of its production and all *E. sibthorpiana* extracts were more effective (10.69%–35.22%) than used controls in this test (Figure 3).

In summary, the lowest activity of studied species was recorded in anti-biofilm assay where ethanol and methanol extracts of both species and essential oil of *E. sibthorpiana* did not show potential for reduction of biofilm synthesis. In contrast, analyzed samples showed the best activity in anti-pyocyanin test, which further enhanced their inhibitory effects on QS mechanism.

Previously, it was published that essential oil of aerial parts of *Ferula asafoetida* (Apiaceae) was active against used virulent factors of *P. aeruginosa*, while essential oil of *Dorema aucheri*




Fig. 2. Light microscopy of colony edges of PAO1 in twitching motility, grown in the presence or absence of *Echinophora sibthorpiana* extracts, essential oil and antibiotics. The bacterial colonies grown with the presence of 0.5 MIC of extracts (A-F) and essential oil (G); *Pseudomonas aeruginosa* colony in the presence of streptomycin had reduced protrusion (H); colony in the presence of ampicillin with regularly formed protrusions (I); control *Pseudomonas aeruginosa* produced a flat, widely spread, irregularly shaped colony (J); Magnification: (A-J) × 100.

did not effect biofilm and pyocyanin synthesis (Sepahi et al., 2015). As previously mentioned, *E. sibthorpiana* essential oil did not show potential in the reduction of biofilm synthesis, while *F. macedonica* oil expressed low activity. In our work, the anti-QS effects of tested extracts varied against different virulent factors, which is in agreement with the study of Adonizio (2008) where various plant extracts had different activity on *P. aeruginosa*. This bacterium uses the ability to form a biofilm controlled by gene expression, as a resistance to the antibiotics and as a protection from the host's immune




Fig. 3. Reduction of pyocyanin production of PAO1 by subMIC of *Ferulago macedonica* and *Echinophora sibthorpiana* samples and antibiotics; ME_ap - methanolic extract aerial part; ME_r - methanolic extract root; EE_ap - ethanolic extract aerial part; EE_r - methanolic extract root; AE_ap - methanolic extract aerial part; AE_r - methanolic extract root; EO_ap - essential oil aerial part; STR - Streptomycin; APC - Ampicillin; Ctrl. - Control PAO1.

system. Anti-biofilm test applied in this study detects the inhibition of initial phase of biofilm formation (Soković et al., 2014) and according to obtained results, our samples demonstrated activity in a dosage-dependent manner. Tested essential oils in this study showed strong inhibition of the cells motility and the growth of the colonies. On the other hand, essential oils of *Apium graveolens*, *Foeniculum vulgare*, *Petroselinum crispum* and *Trachyspermum ammi* (Apiaceae) did not show anti-QS potential on *C. violaceum* CV12472 and CVO26 strains (Khan et al., 2009). Obtained differences in the activity of the samples recorded for anti-biofilm and anti-pyocyanin tests may originate from the complex chemical composition of the extracts and essential oils and the possible synergism of various components (Adonizio, 2008). According to presented data in this work, ethanol extracts of *E. sibthorpiana* roots had moderate anti-QS activity and similar results were observed by Nagy (2010) for *Eryngium aquaticum* and *Foeniculum vulgare* root ethanol extracts. Aqueous extracts of both species tested in this study had lower anti-QS potential. These results may suggest a different polarity of ingredients active in the reduction of PAO1 virulence. The method used for extracts preparation and essential oils isolation also should be taken into consideration in results discussion (Adonizio, 2008).

CONCLUSION

In this work, tested concentrations of extracts and essential oils of *F. macedonica* and *E. sibthorpiana* were mostly sufficient to exhibit an inhibitory effect on selected *P. aeruginosa* determinants. In general, it can be concluded that extracts and oil of *E. sibthorpiana* had higher anti-QS potential. The most prominent were the effects of the samples in inhibition of pyocyanin synthesis, where the strongest were essential oils of both species. According to tested parameters of *P. aeruginosa* cells such as the colors and diameters of the colonies, the presence and diameters of flagella, the strongest tested agent was *E. sibthorpiana* oil. In contrast, used amounts of the samples showed low or lack of efficiency in suppressing of biofilm synthesis. All samples were more effective in anti-pyocyanin assay in comparison to the activity of streptomycin and ampicillin. In addition, few samples had stronger activity in twitching flagella test which supported their anti-QS properties considering intercellular communication of *P. aeruginosa*.

ACKNOWLEDGMENTS

Acknowledgment. The authors are grateful to the Ministry of Education, Science and Technological Development of the Republic of Serbia for financial support (Grants No. 173029 and 173021).

REFERENCES

- Adonizio, A. L. (2008). Anti-quorum sensing agents from south Florida medicinal plants and their attenuation of *Pseudomonas aeruginosa* pathogenicity, *ProQuest ETD Collection for FIU* pp. 1–171.
- Ahmad, V. U., Jassbi, A. R. and Pannahi, M. S. C. (1999). Analysis of the Essential Oil of *Echinophora sibthorpiana* Guss. by Means of GC, GC/MS and ¹³C-NMR Techniques, *Journal of Essential Oil Research* 11(1): 107–108.
- Ali, A., Tabanca, N., Ozek, G., Ozek, T., Aytac, Z., Bernier, U. R., Agramonte, N. M., Baser, K. H. C. and Khan, I. A. (2015). Essential Oils of *Echinophora lamondiana* (Apiaceae: Umbelliferae): A Relationship Between Chemical Profile and Biting Deterrence and Larvicidal Activity Against Mosquitoes (Diptera: Culicidae), *Journal of Medical Entomology* 52(1): 93–100.
- Avijgan, M., Hafizi, M., Saadat, M. and Nilforoushzadeh, M. A. (2010). Antifungal Effect of *Echinophora platyloba*' s Extract against *Candida albicans*, *Iranian Journal of Pharmaceutical Research* 0(0): 285–289.
- Avijgan, M., Mirzadeh, F. and Ahmadi Nia, E. (2012). The comparative study of anti-fungal effect of pharmaceutical products containing hydroalcoholic extract of *Echinophora platyloba* DC and fluconazole in women with chronic recurrent vaginitis caused by candida albicans, *Research Journal of Medical Sciences* 17: S 103–S 107.
- Baser, K. H., Erdemgil, F. Z. and Özek, T. (1994). Essential Oil of *Echinophora tenuifolia* L. subsp. *sibthorpiana* (Guss.) Tutin, *Journal of Essential Oil Research* 6(4): 399–400.
- Baser, K. H., Kürkcüoğlu, M., Malyer, H. and Bıcaçci, A. (1998). Essential Oils of Six *Echinophora* Species from Turkey, *Journal of Essential Oil Research* 10(3): 345–351.
- Bauer, W. D. and Mathesius, U. (2004). Plant responses to bacterial quorum sensing signals, *Current Opinion in Plant Biology* 7(4): 429–433.
- Cakilcioglu, U. and Turkoglu, I. (2010). An ethnobotanical survey of medicinal plants in Sivrice (Elazığ-Turkey), *Journal of Ethnopharmacology* 132(1): 165–175.
- Canonn, J. (1968). *Ferulago* W. Koch., in T. Tutin, V. Heywood, N. Burges, D. Moore, D. Valentine, S. Walters and D. Webb (eds), *Flora Europaea*, 2nd edn, Cambridge University Press, London, UK, p. 359–360.
- Cecchini, C., Coman, M. M., Cresci, A., Tirillini, B., Cristalli, G., Papa, F., Sagratini, G., Vittori, S. and Maggi, F. (2010). Essential oil from fruits and roots of *Ferulago campestris* (Besser) Grecescu (Apiaceae): composition and antioxidant and anti-*Candida* activity, *Flavour and Fragrance Journal* 25(6): 493–502.
- Değirmencioglu, N., Gocmen, D., Dağdelen, A. and Dağdelen, F. (2005). Influence of Tarhana Herb (*Echinophora sibthorpiana*) on Fermentation of Tarhana, Turkish Traditional Fermented Food, *Food Technology and Biotechnology* 43: 175–179.

- Delaram, M., Kheiri, S. and Hodjati, M. R. (2011). Comparing the Effects of *Echinophora platyloba*, Fennel and Placebo on Pre-menstrual Syndrome, *Journal of Reproduction & Infertility* **12**(3): 221–226.
- Demirci, S. and Özkan, E. (2014). Ethnobotanical studies of some apiaceae plants in kahramanmaras and a review of their phytochemical studies, *Journal of Faculty Pharmacy of Istanbul University* **44**(2): 241–250.
- Drenkard, E. and Ausubel, F. M. (2002). Pseudomonas biofilm formation and antibiotic resistance are linked to phenotypic variation, *Nature* **416**(6882): 740–743.
- Eadie, M. (2004). The antiepileptic Materia Medica of Pediacus Dioscorides, *Journal of Clinical Neuroscience* **11**(7): 697–701.
- Entezari, M., Hashemi, M., Ashki, M., Ebrahimian, S., Bayat, M., Saraji, A. and Rohani, S. (2009). Studying the effect *Echinophora platyloba* extract on bacteria (*Staphylococcus aureus* and *Pseudomonas aeruginosa*) and fungi (*Candida albicans*, *Aspergillus flavus* and *Aspergillus niger*) *in vitro*, *World Journal of Medical Science* **4**: 89–90.
- Evergetis, E., Michaelakis, A. and Haroutounian, S. A. (2013). Exploitation of Apiaceae family essential oils as potent biopesticides and rich source of phellandrenes, *Industrial Crops and Products* **41**: 365–370.
- Gholivand, M., Rahimi-Nasrabadi, M., Mehraban, E., Niasari, M. and Batooli, H. (2011). Determination of the chemical composition and *in vitro* antioxidant activities of essential oil and methanol extracts of *Echinophora platyloba* DC, *Natural Product Research* **25**(17): 1585–1595.
- Glamočlija, J., Ćirić, A., Nikolić, M., Fernandes, n., Barros, L., Calhelha, R. C., Ferreira, I. C., Soković, M. and van Griensven, L. J. (2015). Chemical characterization and biological activity of Chaga (*Inonotus obliquus*), a medicinal "mushroom", *Journal of Ethnopharmacology* **162**: 323–332.
- Gokbulut, I., Bilenler, T. and Karabulut, I. (2013). Determination of Chemical Composition, Total Phenolic, Antimicrobial, and Antioxidant Activities of *Echinophora tenuifolia* Essential Oil, *International Journal of Food Properties* **16**(7): 1442–1451.
- Gurbuz, O., Gocmen, D., Ozmen, N. and Dagdelen, F. (2010). Effects of yeast, fermentation time, and preservation methods on tarhana, *Preparative Biochemistry and Biotechnology* **40**(4): 263–275.
- Khalighi-Sigaroodi, F., Hadjiakhoondi, A., Shahverdi, A. R., Mozaffarian, V.-A. and Shafiee, A. (2005). "Chemical composition and antimicrobial activity of the essential oil of *Ferulago bernardii* Tomk. and M. Pimen", *DARU Journal of Pharmaceutical Sciences* **13**(3): 100–104.
- Khan, M., Zahin, M., Hasan, S., Husain, F. and Ahmad, I. (2009). Inhibition of quorum sensing regulated bacterial functions by plant essential oils with special reference to clove oil, *Letters in Applied Microbiology* **49**(3): 354–360.
- Micevski, K. (2005). *Echinophora* L., in V. Matevski (ed.), *Flora of Republic of Macedonia*, 6th edn, Macedonian Academy of Sciences and Arts, Skopje, p. 1543–1544.
- Mileski, K., Džamic, A., Ćirić, A., Grujić, S., Ristić, M., Matevski, V. and Marin, P. (2014). Radical scavenging and antimicrobial activity of essential oil and extracts of *Echinophora sibthorpiana* Guss. from Macedonia, *Archives of Biological Sciences* **66**(1): 401–413.
- Mileski, K., Džamic, A., Ćirić, A., Ristić, M., Grujić, S., Matevski, V. and Marin, P. (2015). Composition, antimicrobial and antioxidant properties of endemic species *Ferulago macedonica* Micevski & E. Mayer, *Records of Natural Products* **9**(2): 208–223.
- O'Toole, G. A. and Kolter, R. (1998a). Initiation of biofilm formation in *Pseudomonas fluorescens* WCS365 proceeds via multiple, convergent signalling pathways: a genetic analysis, *Molecular Microbiology* **28**(3): 449–461.
- O'Toole, G. A. and Kolter, R. (1998b). Flagellar and twitching motility are necessary for *Pseudomonas aeruginosa* biofilm development, *Molecular Microbiology* **30**(2): 295–304.
- Ozdemir, S., Gocmen, D. and Yildirim Kumral, A. (2007). A Traditional Turkish Fermented Cereal Food: Tarhana, *Food Reviews International* **23**(2): 107–121.
- Ozkan, A. M. G., Demirci, B., Demirci, F. and Başer, K. H. C. (2008). Composition and Antimicrobial Activity of Essential Oil of *Ferulago longistylis* Boiss. Fruits, *Journal of Essential Oil Research* **20**(6): 569–573.
- Ph.Eur.8.0. (2013). *European Pharmacopoeia 8.0*, Council of Europe, Strasbourg.
- Razavi, S. M., Ravansalar, A. and Mirinejad, S. (2015). The investigation on phytochemicals from *Ferulago angulata* (Schlecht) Boiss, indigenous to central parts of Iran, *Natural Product Research* **29**(21): 2037–2040.
- Ruberto, G., Cannizzo, S., Amico, V., Bizzini, M. and Piattelli, M. (1994). Chemical Constituents of *Ferulago nodosa*, *Journal of Natural Products* **57**(12): 1731–1733.
- Saei-Dehkordi, S. S., Fallah, A. A., Saei-Dehkordi, S. S. and Kousha, S. (2012). Chemical Composition and Antioxidative Activity of *Echinophora platyloba* DC. Essential Oil, and Its Interaction with Natural Antimicrobials against Food-Borne Pathogens and Spoilage Organisms, *Journal of Food Science* **77**(11): M631–M637.
- Sepahi, E., Tarighi, S., Ahmadi, F. S. and Bagheri, A. (2015). Inhibition of quorum sensing in *Pseudomonas aeruginosa* by two herbal essential oils from Apiaceae family, *Journal of Microbiology* **53**(2): 176–180.
- Sharafati-chaleshtori, R., Rafieian-kopaei, M., Mortezaei, S., Sharafati-chaleshtori, A. and Amini, E. (2012). Antioxidant and antibacterial activity of the extracts of *Echinophora platyloba* D.C, *African Journal of Pharmacy and Pharmacology* **6**: 2692–2695.
- Soković, M., Ćirić, A., Glamočlija, J., Nikolić, M. and van Griensven, L. (2014). Agaricus Blazei Hot Water Extract Shows Anti Quorum Sensing Activity in the Nosocomial Human Pathogen *Pseudomonas aeruginosa*, *Molecules* **19**(4): 4189–4199.
- Spoering, A. L. and Lewis, K. (2001). Biofilms and Planktonic Cells of *Pseudomonas aeruginosa* Have Similar Resistance to Killing by Antimicrobials, *Journal of Bacteriology* **183**(23): 6746–6751.
- Taran, M., Reza Ghasempour, H. and Ebrahim, S. (2010). Antimicrobial activity of essential oils of *Ferulago angulata* subsp. *carduchorum*, *Jundishapur Journal of Microbiology* **3**.
- Teplitski, M., Robinson, J. B. and Bauer, W. D. (2000). Plants Secrete Substances That Mimic Bacterial N -Acyl Homoserine Lactone Signal Activities and Affect Population Density-Dependent Behaviors in Associated Bacteria, *Molecular Plant-Microbe Interactions* **13**(6): 637–648.

- Tutin, T. (1968). *Echinophora* L., in T. Tutin, V. Heywood, N. Burges, D. Moore, D. Valentine, S. Walters and D. Webb (eds), *Flora Europaea*, 2nd edn, Cambridge University Press, London, UK, p. 324.
- Valizadeh, H., Mahmoodi, F., Alizadeh, Z. and Bahadori, M. B. (2014). Isolation and Structure Elucidation of Secondary Metabolites from *Echinophora platyloba* DC from Iran, *Journal of Medicinal Plants* **13**: 15–21.
- Yeo, S. S. M. and Tham, F. Y. (2012). Anti-quorum sensing and antimicrobial activities of some traditional Chinese medicinal plants commonly used in South-East Asia, *Malaysian Journal of Microbiology* **8**(1): 11–20.
- Youse, H. A., Kazemian, A., Sereshti, M., Rahmanikhoh, E., Ahmadiania, E., Razaian, M., Maghsoodi, R. and Darani, H. Y. (2012). Effect of *Echinophora platyloba*, *Stachys lavandulifolia*, and *Eucalyptus camaldulensis* plants on *Trichomonas vaginalis* growth in vitro, *Advanced Biomedical Research* **1**: 79.